

Conservation Laws in Field Dynamics or Why Boundary Motion is Exactly Integrable?

Mark B. Mineev–Weinstein

Center for Nonlinear Studies, MS–B258
Los Alamos National Laboratory
Los Alamos, NM 87545

tel: 1(505)667-9186; fax: 1(505)665-3003;
e-mail: mineev@goshawk.lanl.gov

February 9, 2008

Abstract

An infinite number of conserved quantities in the field dynamics $\phi_t = L U(\phi) + \rho$ for a linear Hermitian (or anti-Hermitian) operator L , an arbitrary function U and a given source ρ are presented. These integrals of motion are the multipole moments of the potential created by ϕ in the far-field. In the singular limit of a bistable scalar field $\phi = \phi_{\pm}$ (i.e. Ising limit) this theory describes a dissipative boundary motion (such as Stefan or Saffman-Taylor problem that is the continuous limit of the DLA-fractal growth) and can be exactly integrable. These conserved quantities are the polynomial conservation laws attributed to the integrability. The criterion for integrability is the uniqueness of the inverse potential problem's solution.

Key words: integrability, boundary motion, phase-field model.

PACS numbers: 47.15. Hg, 68.10.-m, 68.70.+w, 47.20. Hw.

Motivation: Two-dimensional (2D) Laplacian growth (known also as Hele-Shaw, Saffman-Taylor, or the quasistatic Stefan problem) has many attributes of an exactly integrable nonlinear problem [1]. At the same time it is dissipative, which is not usually the case for exactly integrable theories. These problems of moving boundaries belong to the more general class of pattern formation phenomena which have been studied intensively for the last decade [2]. From different first-principles models describing pattern formation we will choose and discuss here the so-called phase-field model [3] derived from the basics of the thermodynamics and describing the kinetics of various phase transitions, such as different kinds of solidification (from the supercooled liquid, from the supersaturated solution, from the electrolyte), phase transformations in liquid crystals and in other condensed matter systems, etc. It is known that this phase-field model in the appropriate singular limit is equivalent to the symmetrical Stefan problem [4]. The following elementary (but unexpected) observation motivated this work: a quite general field dynamics (Eq.(1) below) appears to have an infinite number of conservation laws such that in the singular limit the appropriate interface dynamics can be an exactly integrable (dissipative) theory. Besides its purely fundamental interest, this result can help one to study of cases with nonzero (and even anisotropic) surface tension. It turns out that the conservation laws in the presence of surface tension are not lost, but are only corrected (Eq.(35) below). Thus in the framework of exact integrability one can eliminate the short-wave instability, which is of great physical and practical importance, because without this stabilization the problem of Laplacian growth is ill-posed [2].

Conserved integrals: Let us consider the following dynamics of a field $\phi(x, t)$ on the D -dimensional manifold M ($x \in M$):

$$\phi_t = L U(\phi) + \rho \quad (1)$$

where ϕ_t means partial derivative of the ϕ with respect to time t , L is a linear Hermitian or anti-Hermitian operator, U is an arbitrary (generally nonlinear) function of ϕ (and may also be a function of ϕ_t and of $\nabla\phi$), and $\rho(x, t)$ is a given source term.

Let us consider solutions, $\psi_n(x)$, of the equation

$$L \psi_n = 0 \quad (2)$$

such that the integrals

$$c_n(t) = \int_M \psi_n \phi d^D x \quad (3)$$

are finite.

Here if $D > 1$, the zero eigenvalue of L is infinitely degenerate, where n can be discrete or continuous. In general, n stands for the appropriate set of numbers designating the eigenfunctions of zero-eigenvalue of L . For example, if L is the Laplacian and $D = 2$ we have

$$\psi_n = z^{\pm n} \quad (4)$$

Here $n = 0, 1, 2, \dots, \infty$; and z is a complex coordinate on the plane $z = x + iy$. For $D = 3$,

$$\psi_n^{(m)} = P_{n-1}^{(m)}(\theta) e^{im\omega} / r^n \quad (5)$$

or

$$= P_n^{(m)}(\theta) e^{im\omega} r^n \quad (6)$$

Here, r , θ , and ω are the polar coordinates and $P_l^{(n)}(\theta)$ are the associated Legendre polynomials with $|m| < n$. Also $n > 0$ in Eq.(5).

One can immediately see that the dynamics of the integrals c_n is known and given by

$$\begin{aligned} dc_n/dt &= \int_M \psi_n \phi_t d^D x \\ &= \int_M \psi_n L U(\phi) d^D x + \int_M \psi_n \rho d^D x = \int_M \psi_n \rho d^D x \quad . \end{aligned} \quad (7)$$

The first integral vanishes because L is Hermitian (or anti-Hermitian):

$$\int_M \psi_n L U(\phi) d^D x = \pm \int_M U(\phi) L \psi_n d^D x = 0 \quad . \quad (8)$$

There are two important particular cases:

A) ρ is constant in time. In this case the integrals c_n 's are linear in time.

B) $\rho = 0$. In this case the c_n 's are constants in time.

Hamiltonian and dissipative processes: The dynamics (1) is quite general. Many important physical processes are described by this equation with $U = \delta H / \delta \phi$, where (in the language of the theory of phase transitions) $H[\phi]$, ϕ , and U are the free energy functional, the order parameter, and the external field respectively. We do not specify the nature of ϕ yet: it can be a scalar, vector, tensor, etc. For the $\rho = 0$ case, the corresponding dynamical equation is:

$$\phi_t = L \frac{\delta H}{\delta \phi} \quad (9)$$

and can describe both Hamiltonian and dissipative processes. In the former case H is a Hamiltonian, in the latter one H can be a Lyapunov functional, i.e. a functional decreasing in time.

An example of a Hamiltonian process is:

$$\phi_t = \nabla \times \frac{\delta H}{\delta \phi} \quad (10)$$

Here ϕ is a vector and $L = \mathbf{curl}$. In this case the zero eigenvalue functions are $\psi = \mathbf{grad} W$ with arbitrary W , and it is easy to see that H is constant in time:

$$\frac{dH}{dt} = \int \frac{\delta H}{\delta \phi} \phi_t d^D x = \int \frac{\delta H}{\delta \phi} [\nabla \times \frac{\delta H}{\delta \phi}] d^D x = 0 \quad (11)$$

An example of a dissipative process is:

$$\phi_t = (a\nabla^2 - b) \frac{\delta H}{\delta \phi} \quad (12)$$

Here ϕ is a scalar and $L = a\nabla^2 - b$, where a and b are positive real numbers. It is easy to see that if there is no source ($\rho = 0$), H decreases in time:

$$\begin{aligned} \frac{dH}{dt} &= \int \frac{\delta H}{\delta \phi} \phi_t d^D x = \int \frac{\delta H}{\delta \phi} (a\nabla^2 - b) \frac{\delta H}{\delta \phi} d^D x \\ &= - \int \{a (\nabla \frac{\delta H}{\delta \phi})^2 + b (\frac{\delta H}{\delta \phi})^2\} d^D x < 0 \end{aligned} \quad (13)$$

Mathematical interpretation of the quantities c_n 's: For the particular case where Eq.(1) describes a Hamiltonian process, L is skew-symmetric, and a Poisson bracket $\{, \}$ can be introduced. The integrals c_n 's are Casimirs [5], i.e. such functionals, that commute with all others: $\{c_n, F\} = 0$ for all F . Since we are mainly concerned with *dissipative processes and L in this case, as a rule, is not skew-symmetric*, we do not have such an interpretation of the c_n 's. We can only say that the set of c_n 's determines the projection of the field ϕ onto the null space of the operator L , (i.e. the space spanned by the eigenfunctions ψ_n of L with zero eigenvalue). The knowledge of the c_n 's provides only partial information about ϕ . In the source-free case, the dynamics of ϕ occurs outside the null space while the projection of ϕ onto the null-space of L is constant in time. So a great deal of information concerning the field ϕ is beyond our knowledge if the c_n 's are all that we know.

Physical interpretation of the c_n 's: If L is a second order elliptic operator, the quantities c_n 's are the coefficients of the multipole expansion of the

potential $\Phi(t, x)$ created by a distribution of “matter” with density $\phi' = \phi - \lim_{|x| \rightarrow \infty} \phi$ in the far-field:

$$L \Phi = \phi' \quad (14)$$

To see this, we write the solution of Eq.(14) as

$$\Phi(x) = \int G(x, y) \phi'(y) d^D y \quad (15)$$

where

$$L G(x, y) = \delta(x - y) \quad (16)$$

and expand the Green’s function $G(x, y)$ as

$$G(x, y) = \sum_n \psi'_n(x) \psi_n(y) \quad (17)$$

where ψ'_n and ψ_n are the two linearly independent solutions of the equation $L \psi = 0$ corresponding to a given n . Assuming the commutativity of the sum and the integral, we finally get

$$\Phi(t, x) = \sum_n c_n(t) \psi'_n(x) \quad (18)$$

Let us take the Laplacian cases $L = \nabla^2$ for R^2 and R^3 and assume the field $\phi'(x)$ to be localized so that it is zero in the far-field. In the 2D case

$$\Phi(t, z) = \text{Re} \sum_n c_n(t) z^{-n} \quad (19)$$

where

$$c_n(t) = \int_{R^2} z^n \phi(t; x, y) dx dy, \quad (20)$$

and in the 3D case

$$\Phi(t, x) = \sum_n \sum_m c_{nm}(t) P_n^{(m)}(\theta) e^{im\omega} / r^{(n+1)} \quad (21)$$

here x stands for a vector-position and

$$c_{nm}(t) = \int_{R^3} P_n^{(m)}(\theta) e^{im\omega} r^n \phi(t; r, \theta, \omega) d^3 x \quad (22)$$

Now the question arises: how much can we say about the density of “matter” $\phi(t, x)$ from the knowledge of the potential created by ϕ in the far field? This is a classical inverse potential problem having in the general case an infinity of solutions. One can imagine for example a charged ball with initially a spherically symmetrical but inhomogeneous charge density

$\phi(0, r)$. It is clear that all the c_n 's except c_0 are zero in this case while $\phi(t, r)$ can be distributed arbitrarily. So we have an infinite number of solutions for ϕ satisfying the infinite number of constraints $c_n = c_0 \delta_{0,n}$ where $\delta_{0,n}$ is the Kronecker-symbol, and so the same constants c_n 's can correspond to different fields $\phi(t, x)$ (see ref.[6]).

Stefan problem as a singular limit of the “asymmetric” phase-field dynamics: Now we consider the process described by the following equations:

$$\begin{cases} \phi_t + \beta U_t = \frac{1}{2} \nabla [((D_+ + D_-) + (D_+ - D_-)\phi) \nabla U] \\ \alpha \xi^2 \phi_t = \xi^2 \nabla^2 \phi + (\phi - \phi^3)/a + U \end{cases} \quad (23)$$

This equations encompass several well-known problems. If $D_+ = D_-$ the system (23) coincides with the phase-field model [3] which describes a broad class of pattern formation problems where the order parameter ϕ is called a “phase”. D_+ and D_- stand for the diffusions coefficients in the bulk where $\phi = 1$ and $\phi = -1$ respectively (i.e. far from the “interface” defined as the surface described by $\phi(t, x) = 0$). The first equation in the system (23) is a particular case of Eq.(1) for $\beta = 0$, and the second one is the time-dependent Ginzburg-Landau equation [7] in the presence of external field U . Here $\alpha \xi^2$ and β are kinetic coefficients or the relaxation times of the fields ϕ and U respectively, ξ is the coherence length, and a^{-1} is the double-well potential depth. One can easily see that if $\alpha = \beta = 0$ and $D_+ = D_-$ the system (23) is equivalent to the Cahn-Hilliard equation [8]

$$\phi_t = -\nabla^2(\xi^2 \nabla^2 \phi + (\phi - \phi^3)/a) \quad (24)$$

describing the process of phase separation. It corresponds to the dynamics described by Eq.(9) when $L = \nabla^2$ and the functional H has the Ginzburg-Landau free energy form:

$$H = \int_M \left(\frac{\xi^2 (\nabla \phi)^2}{2} - \frac{\phi^2}{2a} + \frac{\phi^4}{4a} \right) d^D x \quad (25)$$

As it was shown by Caginalp [4] the phase-field model (i.e. the Eqs.(23) with $D_+ = D_-$) in the singular limit $\xi \rightarrow 0$ and $a \rightarrow 0$ reduces to the symmetrical (i.e.when $D_+ = D_-$) Stefan problem

$$\beta U_t = \nabla^2 U \quad x \in M \quad (26a)$$

$$U = \sigma(\kappa + \alpha v_n) \quad x \in \partial\Omega(t) \quad (26b)$$

$$v_n = -[\partial_n U] \quad x \in \partial\Omega(t) \quad (26c)$$

where $\partial\Omega(t)$ is a moving interface between the growing bubble $\Omega(t)$ and the rest of the space, σ is the surface tension proportional to $\xi/\sqrt{a}$, κ is the

mean curvature of the moving boundary $\partial\Omega$, v_n is the normal component of the local velocity of $\partial\Omega(t)$, and $[\partial_n U]$ is the jump of the normal component of the gradient of U , $\partial_n U$, across the boundary $\partial\Omega(t)$. Laplacian growth corresponds to the quasistatic Stefan problem (or Hele-Shaw problem) when $\beta = 0$.

The advantage of the “asymmetric” phase-field model (23) is that the same singular limit ($\xi \rightarrow 0$, $a \rightarrow 0$) reduces the system (23) to the general (not necessarily symmetric) Stefan problem:

$$\beta_{\pm} U_t = \nabla^2 U \quad x \in \Omega_{\pm} \quad (27a)$$

$$U = \sigma(\kappa + \alpha v_n) \quad x \in \partial\Omega \quad (27b)$$

$$v_n = -D_+ \partial_n U|_{phase1} + D_- \partial_n U|_{phase2} \quad x \in \partial\Omega(t) \quad (27c)$$

Here $\beta_{\pm} = \beta/D_{\pm}$, while $\Omega_{\pm}$ are domains corresponding to phases $\phi = \pm 1$.

One can further extend the system (23) to the more general processes described by equations

$$\begin{cases} \phi_t + \beta U_t = (p \nabla q \nabla + r) U \\ \alpha \xi^2 \phi_t = \xi^2 \nabla^2 \phi + F'(\phi)/a + U \end{cases} \quad (28)$$

Here p , q , and r can be functions of ϕ or of the space x . In other words the RHS of the first equation in the (28) is the arbitrary elliptic operator of the second order acting on U . In the RHS of the second equation of (28) the function $F(\phi)$ is introduced such that its derivative with respect to the scalar field ϕ has two stable solutions, $\phi_{\pm}$, and the unstable one, ϕ_0 : $F'(\phi_{\pm}) = F'(\phi_0) = 0$. It is easy to see that the same singular limit as used above reduces the system (28) to the following interface dynamics:

$$\beta_{\pm} U_t = (p_{\pm} \nabla q_{\pm} \nabla + r_{\pm}) U, \quad x \in \Omega_{\pm} \quad (29a)$$

$$U = \sigma(\kappa + \alpha v_n) \quad x \in \partial\Omega(t) \quad (29b)$$

$$v_n = -p[q \partial_n U] \quad x \in \partial\Omega(t) \quad (29c)$$

where $p_{\pm}$, $q_{\pm}$, and $r_{\pm}$ are the functions p , q , and r respectively defined in the domains Ω_+ and Ω_- . It was shown [9] that this kind of interface dynamics (describing rather “elliptic” than Laplacian growth, because of $\nabla^2 \rightarrow p \nabla q \nabla + r$) possesses the infinite number of conservation laws for an arbitrarily dimensional process.

Now we investigate the quantities c_n defined by Eq.(3) for the processes described by (28) in this singular limit:

Conserved integrals c_n 's in the singular limit: The limit $\xi \rightarrow 0$ makes the width between different phases infinitesimally thin, and the limit $a \rightarrow 0$ excludes fluctuations of $\phi(x)$, so it can take only three values: two stable, $\phi_{\pm}$, and one unstable, ϕ_0 . Let us choose for simplicity F in the Ginzburg-Landau form: $F' = \phi - \phi^3$. In this case $\phi_{\pm} = \pm 1$, $\phi_0 = 0$. The equation $\phi_0(t, x) = 0$ describes the boundary moving between two phases $\phi = \pm 1$. By introducing the new function

$$\phi' = (\phi + 1)/2 \quad (30)$$

we describe the interface $\phi(t, x) = 1/2$ moving between two phases described by $\phi' = 1$ in $\Omega(t)$ and $\phi' = 0$ outside it. Consequently, the integrals c_n in this limit are:

$$c_n = \int_{R^D} \psi_n \phi' dx = \int_{\Omega(t)} \psi_n dx \quad (31)$$

But these integrals are the constants of motion found earlier in arbitrarily dimensional boundary motion when $\alpha = \beta = \sigma = 0$ [9]. Thus, in the singular limit $\xi \rightarrow 0$, $a \rightarrow 0$ the conserved quantities, c_n , coincide with the conservation laws previously found. Therefore the field dynamics ((1) is a natural extension of the boundary motion $\partial\Omega(t)$ that possesses an infinite number of conservation laws.

Integrability in the singular limit: An infinite number of polynomial conservation laws is a necessary (but not always sufficient) condition for a nonlinear field problem to be exactly integrable [10]. When do the c_n 's correspond to an integrable case? While the knowledge of the c_n 's is not enough to recover the field $\phi(x)$, the situation is much better in the singular limit described above where the field dynamics $\phi(t)$ is reduced to the boundary dynamics $\partial\Omega(t)$. In this case one might recover the interface and the problem might be solvable. Indeed here the “matter distribution” ϕ' is especially trivial: $\phi' = 1$ in $\Omega(t)$ and $\phi' = 0$ outside it. Therefore looking for the $\phi(t, x)$ by knowing the far-field potential Φ is reduced to the looking for the boundary $\partial\Omega(t)$ between two phases. This problem is called the “inverse potential problem” [11,12] and sometimes can be explicitly solved. For $D = 2$ and $L = \nabla^2$ using conformal mapping it is possible to recover the boundary $\partial\Omega$ from the knowledge of the singularities of the function $\sum_{n=1}^{\infty} c_n z^{-n}$ analytic in R^2/Ω [11], and where c_n are integrals defined by (3). However, if $D \neq 2$ or $L \neq \nabla^2$ we do not have a constructive way to recover the boundary $\partial\Omega$, but if Ω is (a) single-connected and (b) star-like (i.e. a point inside Ω exists such that all straight rays from this point cross the boundary $\partial\Omega$ only once) there exists a unique domain Ω for given potential

[12]. Nevertheless it is still unclear how to parametrize the corresponding boundary in this case. So, although the knowledge of c_n 's in principle is enough to recover the boundary of a single-connected star-like domain in the general multi-dimensional case, only the 2-D Laplacian case is integrable in a constructive way (and not even for all kinds of singularities of the function $\sum_{n=1}^{\infty} c_n z^{-n}$).

Integrals c_n 's in the case of non-zero surface tension $\sigma = \xi/\sqrt{a}$ and non-zero kinetic coefficient α : If in the singular limit $\xi \rightarrow 0$, $a \rightarrow 0$ the ratio $\xi/\sqrt{a}$ has a finite non-zero value that corresponds to a non-zero surface tension, the integrals c_n defined as in Eq.(3) generally speaking are not conserved for the manifold $M = R^D$. (They are even not necessarily finite.) For the process described by the system(23) with $\beta = 0$ the rate of c_m defined by (3) is

$$d c_m / d t = D_- \int_{S^{D-1}} (\psi_m \partial_n U - U \partial_n \psi_m) d S^{D-1} \quad (32)$$

where S^{D-1} is the hypersphere in R^D whose radius tends to infinity when ψ_m diverges at infinity and tends to zero when ψ_m diverges at the origin. (Without surface tension it corresponds to the “internal” and “external” one-side Stefan problems respectively, i.e. when U governed by the equation $LU = 0$ is non-trivial only inside or outside the boundary $\partial\Omega$, while in the complementary phase U vanishes). If $L = \nabla^2$ the function ψ_m diverges (at infinity or at the origin) as an m^{th} power of r (see Eqs.(4)-(6)). In this case, for the RHS of Eq.(32) to vanish, U must decay where ψ_m diverges (i.e. at infinity or at the origin) faster than any power of r . Without surface tension, the field U which is zero in the phase where ψ_m diverges, since U is a harmonic function zero at the boundary, $U = 0$. Thus in this case the RHS of Eq.(32) is zero, so that the c_n 's are conserved.

For non-zero surface tension this is not the case, because U , satisfying non-trivial boundary conditions (see Eq.(27b)) decays at infinity (or at the origin) only algebraically, instead of exponentially (generally speaking as r at the origin or as r^{-1} at the infinity). Therefore, in this case, the RHS of Eq.(32) is not zero, so the c_n 's are not conserved. For the 2D Laplacian internal problem (when $\psi_n = z^n$) it is

$$d c_n / d t = -4 D_- \pi n U_n \quad (33)$$

where

$$U(r, \theta) = \sum_{n=1}^{\infty} (r^{-n} (U_n \exp(-in\theta) + U_{-n} \exp(in\theta))) \quad (34)$$

[13]. By the same reasoning the c_n 's are not conserved when the kinetic coefficient α is not zero.

However, there are situations when $dc_n/dt = 0$ even for non-zero surface tension and kinetic coefficient α : (i) First, this is the case if instead of $L = \nabla^2$ the elliptic operator L is such that its solutions decay exponentially at infinity (for example if $L = \nabla^2 - a$ with a positive a). Evidently the surface integral in the RHS of Eq.(31) also vanishes. (ii) Second, c_n 's are conserved for the *one-side* (i.e. when $D_- = 0$) non-zero surface tension case (which is beyond the phase-field model dealing only with the symmetrical two-side model, when $D_+ = D_-$). The one-side problem with non-zero surface tension is very common for different physical systems, for example for bubbling in Hele-Shaw cell, in solidification when one can neglect diffusion in a solid, etc. [2]. (For simplicity we consider here only isotropic surface tension. For the anisotropic case all the results hold, but σ can no longer be pulled out of integrals and derivatives since it is no longer a number but instead an angle-dependent function.)

It is easy to obtain the rate of change of c_n in this case directly from the boundary motion's formulation (29). Indeed because the integrand in (3) does not depend on time, but only on the domain of integration, we have

$$dc_m/dt = 1/dt \left(\int_{\partial\Omega(t+dt)} - \int_{\partial\Omega(t)} \right) \psi_m d\Gamma$$

(here $d\Gamma$ stands for the element of the moving surface $\partial\Omega(t)$. Further, it equals

$$= \int_{\partial\Omega(t)} v_n \psi_m d\Gamma = \int_{\partial\Omega(t)} \psi_m \partial_n U d\Gamma$$

because of Eq.(29c) (here $p = 1$, $q_+ = 1$, and $q_- = 0$ are chosen for the simplicity);

$$= \int_{\partial\Omega(t)} \psi_m \partial_n U d\Gamma - \int_{\partial\Omega(t)} (U - \sigma(\kappa + \alpha v_n)) \partial_n \psi_m d\Gamma$$

because of Eq.(29b), and finally

$$= -\sigma \int_{\partial\Omega} (\kappa + \alpha \partial_n U) \partial_n \psi_m d\Gamma \quad (36)$$

since the vector field $(U \nabla \psi_m - \psi_m \nabla U)$ is divergenceless because of Eq.(29a):

$$\nabla(U \nabla \psi_m - \psi_m \nabla U) = 0$$

So this is the law of change of c_m defined by (3) for a one-side Stefan model in the presence of surface tension and kinetics on the moving boundary. It is known [14] that without sources of the field U (i.e. when $\rho = 0$)

the boundary $\partial\Omega(t)$ usually has the tendency to take a circular shape in the long-time asymptotics (excluding possible topological changes which are not considered here). In this case we expect $d|c_m^0|/dt$ to be negative since $|c_m^0|$ indicates a deviation from a circular shape. It would be interesting to check this statement in the near future.

Conclusions: We have shown straightforwardly that there exist an infinite number of conserved quantities for the field dynamics described by Eq.(1), such that in the case of a bistable scalar field ϕ the singular limit corresponding to the boundary motion between two phases (Ising limit in the kinetics of phase transitions) can be integrable (for instance if $D = 2$ and $L = \nabla^2$). The criterion of integrability is the solvability of the associated inverse potential problem, where the conserved integrals c_n 's are the coefficients of the multipole expansion of the fictitious potential created by the matter distribution $\phi(x, t)$. In the presence of surface tension and boundary kinetics these conservation laws are broken, but in the framework of the one-side model, the evolution of quantities c_n 's has been calculated (Eq.(35)). It means that in many cases when the integrals c_n 's completely determine the moving cluster, we can follow the interface' evolution with non-zero surface tension.

I am grateful to F. J. Alexander, R. Almgren, and I. M. Gelfand for very helpful discussions of this problem and to S. Ponce-Dawson and P. Sievert for their comments on the paper.

References.

1. S.Richardson *J.Fluid Mech.* **56**, 609 (1972); B.I.Shraiman and D.Bensimon *Phys.Rev.A* **30**, 2840 (1984); D.Bensimon and P.Pelce *Phys.Rev.A* **33**, 4477 (1986); S.D.Howison *J.Fluid Mech.* **167**, 439 (1986); M.B.Mineev *Physica D* **43**, 288 (1990); M.B.Mineev- Weinstein and S.Ponce Dawson *Phys.Rev.E* **50**, R24 (1994); S.Ponce Dawson and M.B.Mineev- Weinstein *Physica D* **73**, 373 (1994)
2. D.Bensimon, L.P.Kadanoff, S.Liang, B.I.Shraiman, and C.Tang *Rev.Mod. Phys.***58**, 977 (1986); *Dynamics of Curved Fronts*, ed. by P.Pelce (Academic Press, San Diego, 1988)
3. J.S.Langer, in *Directions in Condensed Matter Physics*, (World Scientific, Singapore, 1986) p.165
4. G.Caginalp *Phys.Rev. A* **39**, 5887 (1989)
5. B.A.Dubrovin, S.P.Novikoff, and A.T.Fomenko, *Modern Geometry - Methods and Applications*, (Springer - Verlag, New York, 1992)
6. M.B.Mineev-Weinstein and F.J.Alexander *Conserved Moments in Nonequilibrium Field Dynamics*, to be published in Journal of Statistical Physics
7. P.C.Hohenberg and B.I.Halperin, *Rev.Mod.Phys.* **49**, 435 (1977)
8. J.W.Cahn *Acta Metall.* **9**, 795 (1961), J. D. Gunton, M. San Miguel and P. S. Sahni, in *Phase Transitions and Critical Phenomena* ed. by C. Domb and J. L. Lebowitz, Academic, New York, (1983).
9. M.B.Mineev-Weinstein *Phys.Rev.E* **43**, R2241 (1993)
10. M.J.Ablowitz and H.Segur, *Solitons and Inverse Scattering Transform* (SIAM, Philadelphia, 1981)
11. G. Herglotz, *Über die analytische Fortsetzung des Potentials ins Innere der anziehenden Massen*, (Teubner-Verlag, 1914)
12. P.S.Novikoff *Doklady AN SSSR* **18**, 165, (1938)
13. The result in the last paragraph and Eq.(33) were obtained together with R.Almgren.
14. P.Constantin and M.Pugh *Nonlinearity* **6**, 393 (1993)